arXiv:1201.2666v1 [hep-ph] 12 Jan 2012

General Form of the Color Potential Produced by Color Charges of the Quark

Gouranga C Nayak[1,*]

[1] *1001 East 9th Street, #A, Tucson, AZ 85719, USA*

Abstract

Constant electric charge e satisfies the continuity equation $\partial_\mu j^\mu(x) = 0$ where $j^\mu(x)$ is the current density of the electron. However, the Yang-Mills color current density $j^{\mu a}(x)$ of the quark satisfies the equation $D_\mu[A]j^{\mu a}(x) = 0$ which is not a continuity equation ($\partial_\mu j^{\mu a}(x) \neq 0$) which implies that a color charge $q^a(t)$ of the quark is not constant but it is time dependent where $a = 1, 2, ...8$ are color indices. In this paper we derive general form of color potential produced by color charges of the quark. We find that the general form of the color potential produced by the color charges of the quark at rest is given by $\Phi^a(x) = A_0^a(t, \mathbf{x}) = \frac{q^b(t-\frac{r}{c})}{r}\left[\frac{\exp[g\int dr \frac{Q(t-\frac{r}{c})}{r}]-1}{g\int dr \frac{Q(t-\frac{r}{c})}{r}}\right]_{ab}$ where dr integration is an indefinite integration, $Q_{ab}(\tau_0) = f^{abd}q^d(\tau_0)$, $r = |\vec{x} - \vec{X}(\tau_0)|$, $\tau_0 = t - \frac{r}{c}$ is the retarded time, c is the speed of light, $\vec{X}(\tau_0)$ is the position of the quark at the retarded time and the repeated color indices b, d(=1,2,...8) are summed. For constant color charge q^a we reproduce the Coulomb-like potential $\Phi^a(x) = \frac{q^a}{r}$ which is consistent with the Maxwell theory where constant electric charge e produces the Coulomb potential $\Phi(x) = \frac{e}{r}$.

PACS numbers:

---

[*] Electronic address: nayak@max2.physics.sunysb.edu

I. INTRODUCTION

In the eighteenth century, Coulomb, in an impressive series of experiments found that the static potential produced by an electric charge is inversely proportional to distance. This is known as Coulomb's law. The atomic bound state, such as hydrogen atom, is described by using Coulomb potential. However, an exact form of the color potential produced from color charge is not known till now. The exact form of color potential produced by color charges of the quark may provide an insight to the question "why quarks are confined inside a (stable) proton".

Note that, in Maxwell theory fundamental electric charge e of the electron is constant. This constant electric charge e satisfies the continuity equation $\partial_\mu j^\mu(x) = 0$ where $j^\mu(x)$ is the current density of the electron. However, in Yang-Mills theory the Yang-Mills color current density $j^{\mu a}(x)$ of the quark satisfies the equation $D_\mu[A] j^{\mu a}(x) = 0$ which is not the continuity equation ($\partial_\mu j^{\mu a}(x) \neq 0$) where $D_\mu^{ab}[A] = \delta^{ab}\partial_\mu + g f^{acb} A_\mu^c(x)$ and $A^{\mu a}(x)$ is the Yang-Mills potential (color potential). This implies that, unlike electric charge e of the electron which is constant, a color charge of the quark is not constant.

Also, it is important that the conserved color charges are not directly observable – only color representations – because of the unbroken gauge invariance of QCD. Thus, the concept of constant color charge seems unphysical. The asymptotic freedom [1] which includes quantum effects (loop diagrams) predicts that the strong coupling α_s decreases at short distances and increases at long distances.

In quantum mechanics the charge density and probability density of the point particle can be obtained from the quantum wave functions. Hence one finds that the charge density is more microscopic and may depend on the space coordinate $\vec{x}$. For example, consider charge and charge density in a volume $V = \int d^3\vec{x}$. The total charge in the volume $V = \int d^3\vec{x}$ can be obtained from the corresponding charge density after integrating over the entire (physically) allowed volume $V = \int d^3\vec{x}$. Hence one finds that the total charge in the volume $V = \int d^3\vec{x}$ is independent of space coordinate $\vec{x}$. If there is only one point charge in the entire (physically) allowed volume $V = \int d^3\vec{x}$ then one finds that the charge of the point particle is independent of the space coordinate $\vec{x}$.

This implies that if the color charge of the quark is not constant then it has to be time dependent.

Since the color current density $j^{\mu a}(x)$ of a quark has eight color indices $a = 1, 2, ...8$ one finds that a quark has eight time dependent color charges $q^a(t)$ where $a = 1, 2, ...8$. It is useful to remember that the indices $i = 1, 2, 3$ =RED, BLUE, GREEN are not color charges of a quark but they are color indices of the quark field $\psi_i(x)$.

We denote eight time dependent fundamental color charges of a quark by $q^a(t)$ where $a = 1, 2, ...8$ are color indices. A color charge of a quark is flavor independent, *i.e.*, a color charge $q^a(t)$ of the u (up) quark is same as that of the d (down), S (strange), C (charm), B (bottom) or t (top) quark.

It is interesting to note that lattice QCD [2] can not predict the form of the Yang-Mills potential (color potential) $A^a_\mu(x)$ having color indices $a = 1, 2, ..8$ produced from color charge $q^a(t)$. The Wilson loop yields a gauge-invariant measure of the static color singlet quark (effective) potential (energy). In Bohr's atomic model the Coulomb potential provides the force of attraction between proton and electron to form (stable) hydrogen atom. Similarly the color potential (Yang-Mills potential) produced from the color charges may provide the force of attraction between quarks to form (stable) proton.

In the abelian gauge theory the Maxwell equation for electromagnetic potential $A^\mu(x)$ is given by

$$\partial_\mu F^{\mu\nu}(x) = j^\nu(x) \tag{1}$$

where

$$F^{\mu\nu}(x) = \partial^\mu A^\nu(x) - \partial^\nu A^\mu(x). \tag{2}$$

The Dirac equation of the electron is given by

$$[\gamma^\mu \partial_\mu + ie\gamma^\mu A_\mu(x) - m]\psi(x) = 0. \tag{3}$$

The current density of the electron

$$j^\mu(x) = e\bar{\psi}(x)\gamma^\mu\psi(x) \tag{4}$$

obeys the continuity equation

$$\partial_\mu j^\mu(x) = 0. \tag{5}$$

Experimentally, it was discovered by Millikan that the fundamental electric charge e is a constant. From the continuity equation (5) in Maxwell theory we find that the electric charge e is constant which is consistent with the experimental finding of Millikan. Note that we have neglected the vacuum polarization effects. When the vacuum polarization effect is included the effective electric charge e_{eff} increases as r decreases.

The situation in Yang-Mills theory is different, as we will see below.

The Yang-Mills equation for the non-abelian potential $A^{\mu a}(x)$ is given by [3, 4]

$$D_\mu[A]F^{\mu\nu a}(x) = j^{\nu a}(x) \tag{6}$$

where

$$D_\mu^{ab}[A] = \delta^{ab}\partial_\mu + gf^{acb}A_\mu^c(x) \tag{7}$$

and

$$F^{\mu\nu a}(x) = \partial^\mu A^{\nu a}(x) - \partial^\nu A^{\mu a}(x) + gf^{abc}A^{\mu b}(x)A^{\nu c}(x). \tag{8}$$

The Dirac equation of the quark is given by

$$[\delta_{ij}(\gamma^\mu\partial_\mu - m) - igT^a_{ij}\gamma^\mu A^a_\mu(x)]\psi_j(x) = 0. \tag{9}$$

The Yang-Mills color current density of the quark

$$j^{\mu a}(x) = g\bar{\psi}_i(x)T^a_{ij}\gamma^\mu\psi_j(x) \tag{10}$$

obeys the equation

$$D_\mu[A]j^{\mu a}(x) = 0. \tag{11}$$

As discussed above since eq. (11) is not a continuity equation ($\partial_\mu j^{\mu a}(x) \neq 0$), a color charge $q^a(t)$ of the quark is time dependent. For earlier works on classical Yang-Mills theory, see [5–8].

In this paper we derive general form of color potential produced by the color charges of the quark.

Note that the Yang-Mills theory was developed in analogy to the procedure in electromagnetic theory (Maxwell theory) by extending the U(1) gauge group to the SU(3) gauge group appropriately [3]. For example in analogy to $D_\mu\psi = (\partial_\mu - ieA_\mu)\psi$ in electromagnetic theory (Maxwell theory) all derivatives of ψ in Yang-Mills theory appear in the combination $D_\mu\psi = (\partial_\mu - i\epsilon B_\mu)\psi$ where $B_\mu = T^aA^a_\mu$ [3]. Similarly, in analogy to commutator equation

$[D_\mu,\ D_\nu] = ieF_{\mu\nu}$ in electromagnetic theory (Maxwell theory) which predicts the relation between $F_{\mu\nu}(x)$ and $A_\mu(x)$ as given by eq. (2) one finds that in Yang-Mills theory the commutator equation $[D_\mu,\ D_\nu] = -igT^aF^a_{\mu\nu}$ predicts the relation between $F^a_{\mu\nu}(x)$ and $A^a_\mu(x)$ as given by eq. (8) [4]. Similarly, in analogy to the electromagnetic field Lagrangian density $\mathcal{L}_{\rm EM} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$ in electromagnetic theory (Maxwell theory) where $F_{\mu\nu}(x)$ is given by eq. (2) one writes down the Yang-Mills field Lagrangian density $\mathcal{L}_{\rm YM} = -\frac{1}{4}F^a_{\mu\nu}F^{\mu\nu a}$ in Yang-Mills theory where $F^a_{\mu\nu}(x)$ is given by eq. (8) [3]. Hence in order to get an idea of the general form of the color potential (Yang-Mills potential) produced by the color charges of the quark we make analogy between electromagnetic theory (Maxwell theory) and Yang-Mills theory by transforming pure gauge classical potential. The form of the physical electromagnetic potential (Lienard-Wiechert potential or Coulomb potential) produced by the electric charge e of the electron can be obtained from the form of U(1) pure gauge potential produced by the electron. In analogy to the electromagnetic case we find that in Yang-Mills theory the form of physical color potential (Yang-Mills potential) produced by the color charges $q^a(t)$ of the quark can be obtained from the form of SU(3) pure gauge potential produced by the quark.

The above analogy between electromagnetic theory (Maxwell theory) and Yang-Mills theory to obtain the form of the physical potential from the form of the pure gauge potential can be proved by using Lorentz transformation. First of all we note that an electron has non-zero mass (even if very small) and hence it can not travel at speed of light $v = c$. The highest speed of the electron can be $v \sim c$ (which is arbitrarily close to the speed of light). Hence the electron in uniform motion at this highest speed $v \sim c$ produces U(1) (approximate) pure gauge potential. We call this as (approximate) pure gauge potential because the highest speed of the electron can be $v \sim c$ (if $v = c$ then a charge produces exact pure gauge potential). The exact form of the Coulomb potential produced by the electron at rest is obtained from the form of U(1) (approximate) pure gauge potential by putting $v = 0$ in the U(1) (approximate) pure gauge potential. Since the highest speed of the electron can be $v \sim c$ (but not $v = c$), this can also be shown by using Lorentz transformations (see section IV and appendix B). Similarly in Yang-Mills theory a quark has non-zero mass (even if the mass of the light quark is very small) and hence a quark can not travel at speed of light $v = c$. The highest speed of a quark can be $v \sim c$ (which is arbitrarily close to the speed of light). Hence the quark in uniform motion at this highest speed $v \sim c$ produces SU(3) (approximate) pure gauge potential. The exact form of the color potential produced by the quark at rest is obtained from the form of SU(3) (approximate) pure gauge potential by putting $v = 0$ in the SU(3) (approximate) pure gauge potential. Since the highest speed of a quark can be $v \sim c$ (but not $v = c$), this can also be shown by using Lorentz transformations (see section IV).

We find that the general form of the color potential $\Phi^a(x) = A_0^a(t, \mathbf{x})$ produced by the color charges of the quark at rest is given by

$$\Phi^a(x) = A_0^a(t, \mathbf{x}) = \frac{q^b(t - \frac{r}{c})}{r} \left[\frac{\exp[g \int dr \frac{Q(t-\frac{r}{c})}{r}] - 1}{g \int dr \frac{Q(t-\frac{r}{c})}{r}} \right]_{ab} \tag{12}$$

where dr integration is an indefinite integration, a=1,2,..8 are color indices, c is the speed of light, $\tau_0 = t - \frac{r}{c}$ is the retarded time,

$$r = |\vec{x} - \vec{X}(\tau_0)|, \qquad\qquad Q_{ab}(\tau_0) = f^{abd} q^d(\tau_0), \tag{13}$$

$\vec{X}(\tau_0)$ is the position of the quark at the retarded time, and the repeated color indices b, d(=1,2,...8) are summed. The non-zero values of f^{abd} are given by

$$\begin{array}{cccccccccc} \underline{abd} & 123 & 147 & 156 & 246 & 257 & 345 & 367 & 458 & 678 \\ \underline{f^{abd}} & 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} & \frac{\sqrt{3}}{2} & \frac{\sqrt{3}}{2} \end{array} \tag{14}$$

where $f^{abd} = -f^{bad} = -f^{adb}$. All other values of f^{abd} are zero.

For constant color charge q^a, eq. (12) reproduces the Coulomb-like potential $\Phi^a(x) = \frac{q^a}{r}$, which is consistent with the Maxwell theory where constant electric charge e produces Coulomb potential $\Phi(x) = \frac{e}{r}$.

Note that at short distances, the quantum effects via vacuum polarization can be incorporated in to the color charge $q^a(\tau_0)$. The coupling g becomes very small at short distances due to quantum effects/loop diagrams. Hence at short distances where quantum effects are important we find from eq. (12)

$$\Phi^a(x) \to \frac{q^a(\tau_0)}{r}, \qquad \text{as} \qquad g \to 0 \tag{15}$$

which implies that general form of color potential produced by the color charges of the quark reduces to Coulomb-like potential at short distances. This seems to be in agreement with lattice QCD predictions where the effective (color singlet) static confinement (effective) potential energy reduces to Coulomb form at short distances.

Note that the color potential $\Phi^a(x)$ in eq. (12) is not gauge invariant. The gauge invariant quantity is chromo-electric field square $[\sum_{a=1}^8 \vec{E}^a(x)\cdot\vec{E}^a(x)]$. The gauge invariant $[\sum_{a=1}^8 \vec{E}^a(x)\cdot\vec{E}^a(x)]$ can be obtained from eq. (12).

Note that the exact form of color potential produced by color charges of the quark may provide an insight to the question "why quarks are confined inside a (stable) proton". Hence in order to find the exact form of the color potential produced by the color charges of the quark we need to find the exact form of eight time dependent fundamental color charges $q^a(t)$ of the quark where $a = 1, 2, ...8$ are color indices. The derivation of general form of eight time dependent fundamental color charges $q^a(t)$ of a quark in Yang-Mills theory in SU(3) will be presented in a separate publication.

We will present a derivation of eq. (12) in this paper.

The paper is organized as follows. In section II we simplify all the infinite number of non-commuting terms in the SU(3) pure gauge. We obtain general expression of the SU(3) pure gauge potential in section III. In section IV we derive general form of the color potential $\Phi^a(x) = A_0^a(t,\mathbf{x})$ produced by the color charges of the quark at rest as given by eq. (12). Section V contains conclusions.

II. SIMPLIFICATION OF INFINITE NUMBER OF NON-COMMUTING TERMS IN SU(3) PURE GAUGE

The SU(3) pure gauge potential is given by [3, 4]

$$T^aA_\mu^a(x) = \frac{1}{ig}[\partial_\mu U(x)]U^{-1}(x) \tag{16}$$

where T^a is the generator in the fundamental representation of SU(3) group (Gell-Mann matrices) and

$$U(x) = e^{igT^a\omega^a(x)}. \tag{17}$$

The repeated indices a=1,2,...8 are summed. From eq. (17) we find

$$[\partial_\mu U(x)] = [\partial_\mu\omega^d(x)]\frac{d}{d\omega^d(x)}[1 + igT^a\omega^a(x) + \frac{(ig)^2}{2!}T^aT^b\omega^a(x)\omega^b(x) + \frac{(ig)^3}{3!}T^aT^bT^c\omega^a(x)\omega^b(x)\omega^c(x)$$
$$+\frac{(ig)^4}{4!}T^aT^bT^cT^e\omega^a(x)\omega^b(x)\omega^c(x)\omega^e(x) + ...]. \tag{18}$$

Since T^a and T^b do not commute with each other we find from eq. (18)

$$\begin{aligned}
&[\partial_\mu U(x)] = [\partial_\mu \omega^d(x)][igT^d + (ig)^2 T^d T^a \omega^a(x) + \frac{(ig)^3}{2!} T^d T^a T^b \omega^a(x)\omega^b(x) + \frac{(ig)^4}{3!} T^d T^a T^b T^c \omega^a(x) \\
&\omega^b(x)\omega^c(x) + ... \\
&+\frac{(ig)^2}{2!}(T^aT^d - T^dT^a)\omega^a(x) + \frac{(ig)^3}{3!}(T^bT^dT^a + T^aT^bT^d - 2T^dT^aT^b)\omega^a(x)\omega^b(x) \\
&+\frac{(ig)^4}{4!}(T^aT^bT^cT^d + T^bT^cT^dT^a + T^cT^dT^aT^b - 3T^dT^aT^bT^c)\omega^a(x)\omega^b(x)\omega^c(x) + ...].
\end{aligned} \tag{19}$$

Note that the first infinite series in eq. (19) is igT^dU. Hence we find from eq. (19)

$$\begin{aligned}
&[\partial_\mu U(x)] = [\partial_\mu \omega^d(x)][igT^dU + \frac{(ig)^2}{2!}(T^aT^d - T^dT^a)\omega^a(x) + \frac{(ig)^3}{3!}(T^bT^dT^a + T^aT^bT^d - 2T^dT^aT^b) \\
&\omega^a(x)\omega^b(x) + \frac{(ig)^4}{4!}(T^aT^bT^cT^d + T^bT^cT^dT^a + T^cT^dT^aT^b - 3T^dT^aT^bT^c)\omega^a(x)\omega^b(x)\omega^c(x) + ...].
\end{aligned} \tag{20}$$

By using the commutation relation [4]

$$[T^a,\ T^b] = if^{abc}T^c \tag{21}$$

we find from eq. (20)

$$\begin{aligned}
&[\partial_\mu U(x)] = [\partial_\mu \omega^d(x)][igT^dU(x) + \frac{(ig)^2}{2!} if^{ade}\omega^a T^e[1 + igT^b\omega^b(x)] + \frac{(ig)^3}{3!} if^{ade}\omega^a(x) if^{bej}\omega^b(x)T^j \\
&+\frac{(ig)^4}{4!}(if^{ade}T^eT^bT^c + if^{ade}T^bT^eT^c + if^{ade}T^cT^bT^e + if^{ade}T^eT^cT^b + if^{ade}T^bT^eT^c \\
&+if^{ade}T^eT^cT^b)\omega^a(x)\omega^b(x)\omega^c(x) + ...].
\end{aligned} \tag{22}$$

By repeated use of eq. (21) in (22) we finally obtain

$$\begin{aligned}
&[\partial_\mu U(x)] = [\partial_\mu \omega^d(x)][igT^dU(x) + \frac{(ig)^2}{2!} if^{ade}\omega^a(x) T^e[1 + igT^b\omega^b(x)] + \frac{(ig)^3}{3!} if^{ade}\omega^a(x) if^{bej}\omega^b(x)T^j \\
&+\frac{(ig)^4}{4!}(6if^{ade}T^eT^bT^c + 4if^{ade}if^{bej}T^jT^c + if^{ade}if^{bej}if^{cjh}T^h)\omega^a(x)\omega^b(x)\omega^c(x) + ...].
\end{aligned} \tag{23}$$

Simplifying eq. (23) we find

$$\begin{aligned}
&[\partial_\mu U(x)] = [\partial_\mu \omega^d(x)][igT^dU(x) + \frac{(ig)^2}{2!} if^{ade}\omega^a(x) T^e[1 + igT^b\omega^b(x) + \frac{(ig)^2}{2!} T^b\omega^b(x)T^c\omega^c(x)] \\
&+\frac{(ig)^3}{3!} if^{ade}\omega^a(x) if^{bej}\omega^b(x)T^j[1 + igT^c\omega^c(x)] + \frac{(ig)^4}{4!} if^{ade}\omega^a(x) if^{bej}\omega^b(x) if^{cjh}\omega^c(x)T^h + ...].
\end{aligned} \tag{24}$$

Generalizing this to infinity order we obtain

$$\begin{aligned}&[\partial_\mu U(x)] = [\partial_\mu \omega^d(x)][igT^dU(x) + \frac{(ig)^2}{2!}if^{ade}\omega^a(x)T^e[1+igT^b\omega^b(x)+\frac{(ig)^2}{2!}T^b\omega^b(x)T^c\omega^c(x)+....]\\ &+\frac{(ig)^3}{3!}if^{ade}\omega^a(x)if^{bej}\omega^b(x)T^j[1+igT^c\omega^c(x)+....]\\ &+\frac{(ig)^4}{4!}if^{ade}\omega^a(x)if^{bej}\omega^b(x)if^{cjh}\omega^c(x)T^h[1+....]+...].\end{aligned} \tag{25}$$

Note that the expansion in each square bracket is $U(x)$, where $U(x)$ is given by eq. (17). Hence we find from eq. (25)

$$\begin{aligned}&\partial_\mu U(x) = [\partial_\mu \omega^d(x)][igT^dU(x) + \frac{(ig)^2}{2!}if^{ade}\omega^a(x)T^eU(x)+\frac{(ig)^3}{3!}if^{ade}\omega^a(x)if^{bej}\omega^b(x)T^jU(x)\\ &+\frac{(ig)^4}{4!}if^{ade}\omega^a(x)if^{bej}\omega^b(x)if^{cjh}\omega^c(x)T^hU(x)+...].\end{aligned} \tag{26}$$

Multiplying $U^{-1}(x)$ from right in eq. (26) we find

$$\begin{aligned}&[\partial_\mu U(x)]U^{-1}(x) = [\partial_\mu \omega^d(x)][igT^d + \frac{(ig)^2}{2!}if^{ade}\omega^a(x)T^e+\frac{(ig)^3}{3!}if^{ade}\omega^a(x)if^{bej}\omega^b(x)T^j\\ &+\frac{(ig)^4}{4!}if^{ade}\omega^a(x)if^{bej}\omega^b(x)if^{cjh}\omega^c(x)T^h+...].\end{aligned} \tag{27}$$

Simplifying eq. (27) we find

$$\begin{aligned}&\frac{1}{ig}[\partial_\mu U(x)]U^{-1}(x) = [\partial_\mu \omega^b(x)][\delta^{ab} + \frac{ig}{2!}if^{bae}\omega^e(x)+\frac{(ig)^2}{3!}if^{bej}\omega^j(x)if^{ead}\omega^d(x)\\ &+\frac{(ig)^3}{4!}if^{beh}\omega^h(x)if^{ejd}\omega^d(x)if^{jac}\omega^c(x)+...]T^a.\end{aligned} \tag{28}$$

Hence from eqs. (28) and (16) we find

$$\begin{aligned}&T^aA^a_\mu(x) = [\partial_\mu \omega^b(x)][\delta^{ab} + \frac{ig}{2!}if^{bae}\omega^e(x)+\frac{(ig)^2}{3!}if^{bej}\omega^j(x)if^{ead}\omega^d(x)\\ &+\frac{(ig)^3}{4!}if^{beh}\omega^h(x)if^{ejd}\omega^d(x)if^{jac}\omega^c(x)+...]T^a.\end{aligned} \tag{29}$$

From eq. (29) we find that the SU(3) pure gauge potential $A^a_\mu(x)$ is given by

$$A^a_\mu(x) = [\partial_\mu \omega^b(x)]\ [1+\frac{g}{2!}M(x)+\frac{g^2}{3!}M^2(x)+\frac{g^3}{4!}M^3(x)+...]_{ab} \tag{30}$$

where $M_{ab}(x)$ is given by

$$M_{ab}(x) = f^{abc}\omega^c(x). \tag{31}$$

Hence we find from eq. (30) that the SU(3) pure gauge potential is given by

$$A^a_\mu(x) = [\partial_\mu \omega^b(x)]\ [\frac{e^{gM(x)}-1}{gM(x)}]_{ab} \tag{32}$$

where $M_{ab}(x)$ is given by eq. (31).

The abelian-like non-abelian pure gauge is given by

$$\mathcal{A}^a_\mu(x) = \partial_\mu \omega^a(x). \tag{33}$$

It can be seen that, unlike QED, the SU(3) pure gauge in QCD contains infinite number of higher order terms in $\omega^a(x)$. Only the first term in eq. (32) or in eq. (30) corresponds to the U(1) pure gauge in QED, see eq. (33).

III. GENERAL EXPRESSION OF THE SU(3) PURE GAUGE POTENTIAL

Consider the motion of an electron of charge e. Let $X^\mu(\tau)$ be the position of the electron moving with the four-velocity

$$u^\mu(\tau) = \frac{dX^\mu(\tau)}{d\tau}. \tag{34}$$

The Lienard-Wiechert Potential produced by this electron in motion is given by [9]

$$A^\mu(x) = e\frac{u^\mu(\tau_0)}{u(\tau_0)\cdot(x - X(\tau_0))} \tag{35}$$

where x^μ is the time-space position where the electromagnetic field is observed and τ_0 is determined from the solution of the retarded equation

$$x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)|. \tag{36}$$

Writing four-velocity in terms of three-velocity

$$u^\mu = \gamma\beta^\mu = \gamma(\beta_0, \vec{\beta}) = \gamma(1, \vec{\beta}), \qquad \vec{\beta} = \frac{\vec{v}}{c}, \qquad \gamma = \frac{1}{\sqrt{1-\vec{\beta}^2}}. \tag{37}$$

we find from eq. (35) that the Lienard-Weichert potential is given by

$$A^\mu(x) = e\frac{\beta^\mu(\tau_0)}{\beta(\tau_0)\cdot(x - X(\tau_0))}. \tag{38}$$

When $\beta^\mu = (\beta_0, \vec{\beta}) = (1,0,0,0)$ we find from eq. (38)

$$A_0(x) = \frac{e}{x_0 - X_0} = \frac{e}{|\vec{x} - \vec{X}|} \tag{39}$$

which reproduces the Coulomb potential.

Note that since an electron has non-zero mass (even if negligibly small), it can not travel exactly at speed of light $v = c$ but its highest speed can be arbitrarily close to the speed of the light $v \sim c$. On the other hand gluon which carries color charge is massless and travels exactly at speed of light. Hence for a massless particle the four velocity vector β_c^μ is exactly at the speed of light and it remains always at the speed of light where $\beta_c^2 = 0$. Since a quark has non-zero mass (even if the light quark mass is very small), it can not travel exactly at speed of light $v = c$ but the highest speed of the quark can be arbitrarily close to the speed of light $v \sim c$.

For the highest speed of the electron (which is arbitrarily close to the speed of light $v \sim c$) we write

$$\beta_{\sim c}^\mu = (1,\ \vec{\beta}_{\sim c}), \qquad \vec{\beta}_{\sim c}^2 = \frac{\vec{v}^2}{c^2} \sim 1, \qquad \beta_{\sim c}^2 \sim 0 \qquad \text{but} \qquad \beta_{\sim c}^2 \neq 0. \tag{40}$$

From eq. (40) we find

$$u_{\sim c}^\mu = \gamma_{\sim c} \times \beta_{\sim c}^\mu, \qquad \gamma_{\sim c} = \frac{1}{\sqrt{1-\vec{\beta}_{\sim c}^2}} \sim \infty, \qquad \text{but} \qquad \gamma_{\sim c} = \frac{1}{\sqrt{1-\vec{\beta}_{\sim c}^2}} \neq \infty. \tag{41}$$

Note that since electron has non-zero mass we find (see eq. (41))

$$u_{\sim c}^\mu \neq (\infty,\ \infty,\ \infty,\ \infty). \tag{42}$$

For constant speed we find from eq. (35)

$$A^\mu(x) = e\frac{u^\mu}{u \cdot (x - X(\tau_0))}. \tag{43}$$

From eqs. (43), (41) and (40) we find that the electron in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) produces electromagnetic potential

$$A^\mu(x) = e\frac{u_{\sim c}^\mu}{u_{\sim c} \cdot (x - X(\tau_0))} = e\frac{\beta_{\sim c}^\mu}{\beta_{\sim c} \cdot (x - X(\tau_0))}. \tag{44}$$

For electron in uniform motion and at its highest speed $v \sim c$ we find

$$\begin{aligned}\partial^\mu\ e\ln[\beta_{\sim c} \cdot (x - X(\tau_0))] &= e\frac{1}{\beta_{\sim c} \cdot (x - X(\tau_0))}[\beta_{\sim c}^\mu - \beta_{\sim c}^2 \partial^\mu \tau_0] \\ &= e\frac{1}{\beta_{\sim c} \cdot (x - X(\tau_0))}[\beta_{\sim c}^\mu - \beta_{\sim c}^2 \frac{(x - X(\tau_0))^\mu}{u_{\sim c} \cdot (x - X(\tau_0))}] \ \sim\ e\frac{\beta_{\sim c}^\mu}{\beta_{\sim c} \cdot (x - X(\tau_0))}.\end{aligned} \tag{45}$$

From eqs. (40), (41) and (45) we find that eq. (44) can be written as an (approximate) pure gauge potential

$$A^\mu(x) \sim \partial^\mu \omega(x), \qquad \text{where} \qquad \omega(x) = e\ \ln[\beta_{\sim c} \cdot (x - X(\tau_0))]. \tag{46}$$

We call the expression in eq. (46) as (approximate) pure gauge potential because the electron has non-zero mass and hence it can not travel exactly at speed of light $v = c$ or $\beta_c^2 = 0$ to produce the exact pure gauge potential

$$A^\mu(x) = \partial^\mu \omega(x), \qquad \text{where} \qquad \omega(x) = e\ \ln[\beta_{\sim c} \cdot (x - X(\tau_0))]. \tag{47}$$

Hence from eqs. (40), (41), (45), (44) and (46) we find that the expression of the U(1) (approximate) pure gauge potential is given by

$$A^\mu(x) = e\frac{\beta^\mu_{\sim c}}{\beta_{\sim c} \cdot (x - X(\tau_0))} \sim \partial^\mu \omega(x), \qquad \omega(x) = e\ \ln[\beta_{\sim c} \cdot (x - X(\tau_0))] \tag{48}$$

which is the electromagnetic potential produced by the electron of charge e moving at its highest speed (which is arbitrarily close to the speed of the light $v \sim c$).

A. Abelian-Like Pure Gauge Potential and Constant Color Charge

Let us proceed to the non-abelian gauge theory. From eq. (48) we find that the abelian-like (approximate) pure gauge potential produced from the constant color charge q^a is given by

$$\mathcal{A}^a_\mu(x) = q^a\frac{\beta^\mu_{\sim c}}{\beta_{\sim c} \cdot (x - X(\tau_0))} \sim \partial_\mu \omega^a(x), \qquad \omega^a(x) = q^a\ \ln[\beta_{\sim} c \cdot (x - X(\tau_0))] \tag{49}$$

which is the abelian-like potential $\mathcal{A}^a_\mu(x)$ produced by the constant color charge q^a moving at the highest speed (which is arbitrarily close to the speed of the light $v \sim c$). The derivation of eq. (49) is given in appendix A, see eq. (A20).

By using $\omega^a(x)$ from eq. (49) in eq. (32) we find that the non-abelian SU(3) pure gauge potential

$$A^a_\mu(x) = [\partial_\mu \omega^b(x)]\ [\frac{e^{gM(x)} - 1}{gM(x)}]_{ab} = [\partial_\mu \omega^b(x)]\ [1 + \frac{g}{2!}M(x) + \frac{g^2}{3!}M^2(x) + \frac{g^3}{4!}M^3(x) + ...]_{ab} = \partial_\mu \omega^a(x) \tag{50}$$

becomes an abelian-like pure gauge potential $\partial_\mu \omega^a(x)$ where $M_{ab}(x)$ is given by eq. (31). Note that the repeated color indices b(=1,2,....8) are summed in eq. (50).

Hence we find from eq. (50) that if color charge q^a is constant then the SU(3) pure gauge potential $A_\mu^a(x)$ obtained from the SU(3) pure gauge equation $T^a A_\mu^a(x) = \frac{1}{ig}[\partial_\mu U(x)]U^{-1}(x)$ reduces to an abelian-like pure gauge potential $A_\mu^a(x) = \partial_\mu \omega^a(x)$ where $U(x) = e^{igT^a\omega^a(x)}$.

This is expected because when the color charge q^a is constant, the entire physics becomes similar to QED except that the role of constant electric charge e is replaced by constant color charge q^a, where a=1,2,...8. This is equivalent to an abelian-like continuity equation

$$\partial_\mu j^{\mu a}(x) = 0 \tag{51}$$

which becomes possible only when all the generators T^a are diagonal *i.e.*, when $f^{abc} = 0$ for all values of a, b, c=1,2,...8.

However, since all the the generators T^a in SU(3) group are not diagonal the color current density $j_\mu^a(x)$ does not satisfy the continuity eq. (51). Instead, the color current density of the quark satisfies the equation

$$D_\mu[A] j^{\mu a}(x) = 0, \qquad \text{where} \qquad D_\mu^{ab}[A] = \delta^{ab}\partial_\mu + g f^{acb} A_\mu^c(x) \tag{52}$$

which implies that the color charge is not constant.

As discussed in the introduction since the color charge is not constant and it is independent of space coordinate $\vec{x}$ one finds that the color charge has to be time dependent.

Hence we find that unlike constant color charge q^a which does not satisfy eq. (52), the time dependent color charge $q^a(t)$ satisfies eq. (52).

B. General Expression of the SU(3) Pure Gauge Potential

Extending eq. (49) to time dependent color charge $q^a(t)$ we find that the abelian-like non-abelian (approximate) pure gauge potential produced from time dependent color charge $q^a(t)$ is given by

$$\mathcal{A}_\mu^a(x) = q^a(\tau_0) \frac{\beta_{\sim c}^\mu}{\beta_{\sim c} \cdot (x - X(\tau_0))} \sim \partial_\mu \omega^a(x), \qquad \omega^a(x) = \int dl_c \, \frac{q^a(\tau_0)}{l_c}, \qquad l_c = \beta_{\sim c} \cdot (x - X(\tau_0)), \tag{53}$$

which is the abelian-like non-abelian potential $\mathcal{A}_\mu^a(x)$ produced by the time dependent color charge $q^a(t)$ of the quark moving at its highest speed (which is arbitrarily close to the speed of the light $v \sim c$) where dl_c integration is an indefinite integration and τ_0 is obtained from the retarded condition

$$x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)|. \tag{54}$$

The derivation of eq. (53) is given in appendix A, see eq. (A45).

As mentioned earlier a quark has non-zero mass (even if the mass of the light quark is small) which implies that a quark can not travel exactly at the speed of light $v = c$. Hence the highest speed of a quark can be arbitrarily close to the speed of the light $v \sim c$. Hence one finds that a quark in uniform motion at its highest speed (which is arbitrarily close to the speed of the light $v \sim c$) produces (approximate) pure gauge potential.

Note that when color charge q^a is constant we call $\partial_\mu \omega^a(x)$ in eq. (49) as abelian-like pure gauge potential and when color charge $q^a(t)$ is time dependent we call $\partial_\mu \omega^a(x)$ in eq. (53) as abelian-like non-abelian pure gauge potential.

As mentioned above, one can view eq. (49) as the abelian-like (approximate) pure gauge potential which is produced only when all the generators T^a are diagonal. It is similar to soft/collinear photon field. The abelian-like (approximate) pure gauge potential in eq. (49) interacts with quarks but not with gluons even if it carries color indices. On the other hand the abelian-like non-abelian (approximate) pure gauge potential in eq. (53) can interact with gluons as well as with quarks. The soft/collinear gluon field can be obtained from eq. (53).

Using the expressions of $\omega^a(x)$ and $\partial^\mu \omega^a(x)$ from eq. (53) we find from eq. (32) that the general expression of the non-abelian SU(3) (approximate) pure gauge potential is given by

$$A^{\mu a}(x) = q^b(\tau_0) \frac{\beta^\mu_{\sim c}}{\beta_{\sim c} \cdot (x - X(\tau_0))} \left[\frac{\exp[g \int dl_c \frac{Q(\tau_0)}{l_c}] - 1}{g \int dl_c \frac{Q(\tau_0)}{l_c}} \right]_{ab} \tag{55}$$

which is the non-abelian SU(3) Yang-Mills potential (color potential) $A^{\mu a}(x)$ produced by the color charges of the quark in uniform motion at its highest speed (which is arbitrarily close to the speed of the light) where dl_c integration is an indefinite integration and

$$l_c = \beta_{\sim c} \cdot (x - X(\tau_0)), \qquad\qquad Q_{ab}(\tau_0) = f^{abd} q^d(\tau_0). \tag{56}$$

In eqs. (55) and (56) the repeated indices $b, d (= 1, 2, ...8)$ are summed and τ_0 is determined from the solution of the retarded equation

$$x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)|. \tag{57}$$

IV. GENERAL FORM OF COLOR POTENTIAL PRODUCED BY COLOR CHARGES OF THE QUARK

In Maxwell theory we find from eq. (39) that the Coulomb potential is given by

$$A_0(x) = e\frac{\beta_0}{\beta_0(x_0 - X_0)}, \qquad\qquad x_0 - X_0 = |\vec{x} - \vec{X}|, \tag{58}$$

and from eq. (38) we find that the Lienard-Weichert potential is given by

$$A^\mu(x) = e\frac{\beta^\mu(\tau_0)}{\beta(\tau_0)\cdot(x - X(\tau_0))}, \qquad\qquad x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)|, \tag{59}$$

and from eq. (48) we find that the U(1) (approximate) pure gauge potential is given by

$$A^\mu(x) = e\frac{\beta^\mu_{\sim c}}{\beta_{\sim c}\cdot(x - X(\tau_0))}, \qquad\qquad x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)|, \tag{60}$$

where β_0 is the zeroth component of the four velocity vector, $\beta^\mu_{\sim c}$ is the four-velocity arbitrarily close to the speed of light ($v \sim c$) and $\beta^\mu(\tau)$ is any arbitrary four velocity of the electron.

Hence eqs. (58), (59) and (60) suggest that in Maxwell theory (or in U(1) gauge theory) there is just one formula for the electromagnetic potential in the nature. Depending on three different speed limits we call them three different potentials. For example, when speed is zero ($\beta^\mu = (\beta_0,\ 0,\ 0,\ 0)$) we call it Coulomb potential (see eq. (58)), when speed is finite $\Big(\beta^\mu(\tau) = (\beta_0,\ \vec{\beta}(\tau))\Big)$ we call it Lienard-Weichert potential (see eq. (59)) and when speed is arbitrarily close to the speed of light $\Big(\beta^\mu = (\beta_0,\ \vec{\beta}_{\sim c})\Big)$ we call it (approximate) pure gauge potential (see eq. (60)). Eqs. (58), (59) and (60) suggest that if we know one formula then we can get the other two formulas by changing the four-velocity vector.

For example, the static Coulomb potential $A_0(x)$ in eq. (58) can be obtained from the Lienard-Weichert potential $A_\mu(x)$ by replacing

$$\beta^\mu(\tau) \to (\beta_0, 0, 0, 0), \qquad\qquad \text{or by putting} \qquad\qquad \vec{\beta} = 0 \tag{61}$$

in eq. (59). This is obvious because when we put velocity $\vec{\beta} = 0$ then the particle comes to rest, see eq. (37). Similarly the static Coulomb potential $A_0(x)$ in eq. (58) can be obtained from the U(1) (approximate) pure gauge potential $A^\mu(x)$ by replacing

$$\beta^\mu_{\sim c} \to (\beta_0, 0, 0, 0) \tag{62}$$

in eq. (60). Note that $\beta_0 = 1$ remains unchanged in $\beta^\mu(\tau)$ and in $\beta^\mu_{\sim c}$, see eqs. (37) and (40). Since Yang-Mills theory is obtained from Maxwell theory by extending the U(1) gauge group to SU(3) gauge group, one expects eqs. (61) and (62) to be valid in Yang-Mills theory. Hence one expects that the exact form of the Yang-Mills potential produced by the quark moving at any speed can be obtained from the form of the SU(3) (approximate) pure gauge potential produced by the quark in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$).

In other words, one can obtain the exact form of the Yang-Mills potential (color potential) $A_0^a(x)$ produced in a frame where the quark is at rest from the SU(3) (approximate) pure gauge potential $A^{\mu a}(x)$ produced in a frame where the quark is in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) by using eq. (62) in (55).

The above argument can be generalized to a frame where the charge is moving at any arbitrary speed. For example, by comparing eq. (60) with eq. (59) we find that the exact expression of the physical potential (Lienard-Weichert potential) in eq. (59) in Maxwell theory can be obtained from the U(1) (approximate) pure gauge potential by replacing

$$\beta^\mu_{\sim c} \to \beta^\mu(\tau) \tag{63}$$

in eq. (60). This is expected because (approximate) pure gauge potential is obtained from the Lienard-Wiechert potential when the speed of the electron in uniform motion approaches its highest speed (which is arbitrarily close to the speed of light $v \sim c$). By the word "physical potential" we mean the potential which gives physical (gauge invariant) electromagnetic field $F_{\mu\nu}$ in Maxwell theory (see eq. (2)) and physical (gauge invariant) $\sum_{a=1}^{8} F^a_{\mu\nu} F^{\mu\nu a}$ (see eq. (8)) in Yang-Mills theory.

The replacement as given by eq. (62) can also be explicitly verified by using Lorentz transformations. The Lorentz transformation of the electromagnetic potential $A^\mu(x)$ is given by [9]

$$A'^\mu = \frac{\partial x'^\mu}{\partial x^\nu} A^\nu(x), \qquad\qquad A'_\mu = \frac{\partial x^\nu}{\partial x'^\mu} A_\nu(x). \tag{64}$$

Since an electron has non-zero mass it can not travel at speed

$$v = c \tag{65}$$

where c is the speed of light but the highest speed of the electron can be arbitrarily close to the speed of light, *i. e.*,

$$v \sim c. \tag{66}$$

From eqs. (60) we find that an electron in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) produces the electromagnetic potential

$$A^\mu(x) = e\frac{u^\mu_{\sim c}}{u_{\sim c}\cdot(x - X(\tau_0))}. \tag{67}$$

Note that eq. (67) is the U(1) (approximate) pure gauge potential, see eq. (48).

Consider two inertial reference frames K and K' with a relative velocity $\vec{v}$ between them. The time and space coordinates of a point are $(t,\ x,\ y,\ z)$ and $(t',\ x',\ y',\ z')$ in the frames K and K' respectively. If the axes in two inertial frames K and K' remain parallel, but the uniform velocity $\vec{v}$ of the frame K' in frame K is in an arbitrary direction we find [9]

$$x'_0 = u\cdot x, \qquad x'^i = x^i + \frac{\gamma-1}{\vec{\beta}^2}(\vec{\beta}\cdot\vec{x})\beta^i - \gamma\beta^i x_0, \qquad i = 1,2,3. \tag{68}$$

From eqs. (68) and (40) we find

$$x'_0 = u_{\sim c}\cdot x, \qquad x'^i = x^i + \frac{\gamma_{\sim c}-1}{\vec{\beta}^2_{\sim c}}(\vec{\beta}_{\sim c}\cdot\vec{x})\beta^i_{\sim c} - \gamma_{\sim c}\beta^i_{\sim c} x_0, \qquad i = 1,2,3. \tag{69}$$

Hence from eqs. (42) and (69) we find

$$x'_0 \neq \infty, \qquad x'^i \neq \infty, \qquad \text{when} \qquad v \sim c. \tag{70}$$

Since x'_0 and x'^i are not exactly infinity (see eq. (70)) and since the electromagnetic potential $A^\mu(x)$ in eq. (67) is neither infinity nor zero we can perform Lorentz transformation on $A^\mu(x)$ in eq. (67).

From eq. (64) we find

$$A'^a_0 = \frac{\partial x'_0}{\partial x^\nu}A^{\nu a}(x). \tag{71}$$

From eqs. (71), (69) and (67) we find

$$A'_0 = \frac{\partial(u_{\sim c}\cdot x)}{\partial x^\nu}A^\nu(x) = u_{\sim c}\cdot A(x) = e\frac{1}{u_{\sim c}\cdot(x - X(\tau_0))}. \tag{72}$$

From eq. (42) we find that $A'_0 \neq 0$ in eq. (72). Since

$$\tau = \tau' \tag{73}$$

under a Lorentz transformation, we find from eqs. (69) and (72) that

$$A'_0 = e\frac{1}{x'_0 - X'_0(\tau'_0)}. \tag{74}$$

By using the retarded condition

$$x'_0 - X'_0(\tau'_0) = |\vec{x}' - \vec{X}'(\tau'_0)| \tag{75}$$

we find from eq. (74)

$$A'_0 = e\frac{1}{|\vec{x}' - \vec{X}'(\tau'_0)|}. \tag{76}$$

Hence we find from eqs. (73) and (76)

$$A_0 = e\frac{1}{|\vec{x} - \vec{X}(\tau_0)|} \tag{77}$$

which reproduces the exact form of Coulomb potential produced by the electron at rest.

Hence we find that the exact form of the Coulomb potential in eq. (77) can be obtained from the U(1) (approximate) pure gauge potential in eq. (67) by using Lorentz transformations. Hence by using Lorentz transformations we have proved that the static Coulomb potential $A_0(x)$ in eq. (58) can be obtained from the U(1) (approximate) pure gauge potential $A^\mu(x)$ in eq. (60) by using eq. (62) in (60).

From eq. (77) we find that the general form of the electromagnetic potential $A^\mu(x)$ produced by the electron moving with arbitrary four velocity $u^\mu(\tau)$ is given by

$$A_\mu(x) = e\frac{u_\mu(\tau_0)}{u(\tau_0)\cdot(x - X(\tau_0))} \tag{78}$$

which reproduces eq. (59) for the Lienard-Wiechert potential produced by the electron moving with arbitrary four-velocity $u^\mu(\tau)$ where τ_0 is obtained from the solution of the retarded condition

$$x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)|. \tag{79}$$

Similarly one can perform the Lorentz transformation on the electromagnetic field $F^{\mu\nu}(x)$ produced by the electron in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) to obtain the exact form of the static Coulomb electric field produced by the electron at rest (see appendix B).

Hence one finds that once the form of the (approximate) pure gauge electromagnetic potential is known then one can obtain the exact form of the static Coulomb potential from it by using Lorentz transformations. This is consistent with the fact that eq. (58) can be obtained from eq. (60) by using eq. (62) in (60).

Note that the above Lorentz transformation is performed in a frame which is moving arbitrarily close to the speed of light ($v \sim c$) but not exactly at the speed of light ($v = c$). Since the Lorentz transformation on the light-cone remains on the light-cone, one should not make Lorentz transformation at speed of light ($v = c$). However, as shown above, one can make Lorentz transformation in a frame which is moving arbitrarily close to the speed of light ($v \sim c$). This is consistent with the fact that the electron has non-zero mass.

The above Lorentz transformation techniques can be extended to Yang-Mills theory. The Lorentz transformation on the Yang-Mills potential $A^{\mu a}(x)$ is given by [9]

$$A'^{\mu a} = \frac{\partial x'^{\mu}}{\partial x^{\nu}} A^{\nu a}(x), \qquad\qquad A'^{a}_{\mu}(x') = \frac{\partial x^{\nu}}{\partial x'^{\mu}} A^{a}_{\nu}(x). \tag{80}$$

Since a quark has non-zero mass it can not travel at speed

$$v = c \tag{81}$$

where c is the speed of light but the highest speed of the quark can be arbitrarily close to the speed of light, *i. e.*,

$$v \sim c. \tag{82}$$

From eqs. (55), (56) and (57) we find that a quark in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) produces the Yang-Mills potential (color potential)

$$A^{\mu a}(x) = q^{b}(\tau_0) \frac{u^{\mu}_{\sim c}}{u_{\sim c} \cdot (x - X(\tau_0))} \left[\frac{\exp[g \int dl_c \frac{Q(\tau_0)}{l_c}] - 1}{g \int dl_c \frac{Q(\tau_0)}{l_c}}\right]_{ab} \tag{83}$$

where dl_c integration is an indefinite integration and

$$l_c = u_{\sim c} \cdot (x - X(\tau_0)), \qquad\qquad Q_{ab}(\tau_0) = f^{abd} q^{d}(\tau_0). \tag{84}$$

Note that eq. (83) is the SU(3) (approximate) pure gauge potential

$$T^a A^{\mu a}(x) \sim \frac{1}{ig}[\partial^\mu U(x)]U^{-1}(x), \qquad\qquad U(x) = e^{igT^a\omega^a(x)} \tag{85}$$

where (see eq. (53))

$$\omega^a(x) = \int dl_c \,\frac{q^a(\tau_0)}{l_c}. \tag{86}$$

Since a quark has non-zero mass one finds from eq. (70) that x'_0 and x'^i are not exactly infinity. Since x'_0 and x'^i are not exactly infinity and the Yang-Mills potential $A^{\mu a}(x)$ in eq. (83) is neither infinity nor zero we can perform Lorentz transformation on $A^{\mu a}(x)$ in eq. (83).

From eq. (80) we find

$$A'^a_0 = \frac{\partial x'_0}{\partial x^\nu} A^{\nu a}(x). \tag{87}$$

From eqs. (87), (69) and (83) we find

$$A'^a_0 = \frac{\partial(u_{\sim c}\cdot x)}{\partial x^\nu} A^{\nu a}(x) = u_{\sim c}\cdot A^a(x) = \frac{q^b(\tau_0)}{u_{\sim c}\cdot(x - X(\tau_0))}\left[\frac{\exp[g\int dl_c \frac{Q(\tau_0)}{l_c}] - 1}{g\int dl_c \frac{Q(\tau_0)}{l_c}}\right]_{ab}. \tag{88}$$

Since a quark has non-zero mass we find from eq. (42) that $A'^a_0 \neq 0$ in eq. (88). Hence we find from eqs. (69), (73) and (88) that

$$A'^a_0 = \frac{q^b(\tau'_0)}{x'_0 - X'_0(\tau'_0)}\left[\frac{\exp[g\int dl'_0 \frac{Q(\tau'_0)}{l'_0}] - 1}{g\int dl'_0 \frac{Q(\tau'_0)}{l'_0}}\right]_{ab} \tag{89}$$

where

$$l'_0 = x'_0 - X'_0(\tau'_0). \tag{90}$$

By using the retarded condition

$$x'_0 - X'_0(\tau'_0) = |\vec{x}' - \vec{X}'(\tau'_0)| \tag{91}$$

we find from eq. (89)

$$A'^a_0 = \frac{q^b(\tau'_0)}{|\vec{x}' - \vec{X}'(\tau'_0)|}\left[\frac{\exp[g\int dl' \frac{Q(\tau'_0)}{l'}] - 1}{g\int dl' \frac{Q(\tau'_0)}{l'}}\right]_{ab} \tag{92}$$

where

$$l' = |\vec{x}' - \vec{X}'(\tau'_0)|. \tag{93}$$

Hence we find from eqs. (73) and (92)

$$A_0^a = \frac{q^b(\tau_0)}{r}\left[\frac{\exp[g\int dr \frac{Q(\tau_0)}{r}]-1}{g\int dr\frac{Q(\tau_0)}{r}}\right]_{ab} \tag{94}$$

which is the general form of the color potential (Yang-Mills potential) produced by the color charges of the quark at rest where dr is an indefinite integration, $\vec{X}(\tau_0)$ is the position vector of the quark at the retarded time and

$$r = |\vec{x}-\vec{X}(\tau_0)| = x_0 - X_0(\tau_0), \qquad\qquad Q_{ab}(\tau_0) = f^{abd}q^d(\tau_0). \tag{95}$$

In eqs. (94) and (95) the repeated indices $b, d = (1, 2, ...8)$ are summed.

As we saw earlier in Maxwell theory the electric potential (Coulomb potential) in eq. (58) produced by the electron at rest can be obtained from the U(1) (approximate) pure gauge potential in eq. (60) produced by the electron moving at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) by using eq. (62) in (60). Similarly in Yang-Mills theory we find that the color potential in eq. (94) produced by the color charges of the quark at rest can be obtained from the SU(3) (approximate) pure gauge potential in eq. (55) produced by the quark moving at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) by using eq. (62) in (55).

From eq. (94) we find that the general form of the color potential (Yang-Mills potential) $A^{\mu a}(x)$ produced by the color charges of the quark moving with arbitrary four velocity $u^\mu(\tau)$ is given by

$$A_\mu^a(x) = q^b(\tau_0)\frac{u_\mu(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))}\left[\frac{\exp[g\int dl\frac{Q(\tau_0)}{l}]-1}{g\int dl\frac{Q(\tau_0)}{l}}\right]_{ab} \tag{96}$$

where dl integration is an indefinite integration,

$$l = u(\tau_0)\cdot(x-X(\tau_0)), \qquad\qquad Q_{ab}(\tau_0) = f^{abc}q^c(\tau_0), \tag{97}$$

the repeated color indices b, c(=1,2,...8) are summed and τ_0 is determined from the solution of the retarded equation

$$x_0 - X_0(\tau_0) = |\vec{x}-\vec{X}(\tau_0)|. \tag{98}$$

Note that unlike Maxwell theory, the Yang-Mills color current density generated by the Yang-Mills potential in eq. (96) is non-linear function of color charges $q^a(\tau)$, see appendix C.

From the expression of the proper time

$$\tau = \int \frac{dX_0(\tau)}{\gamma(X_0(\tau))} \tag{99}$$

we find that when $\vec{\beta} = 0$

$$\tau_0 = X_0(\tau_0). \tag{100}$$

From the retarded eq. (98) we find

$$x_0 - X_0(\tau_0) = \frac{r}{c} \tag{101}$$

where

$$r = |\vec{x} - \vec{X}(\tau_0)|. \tag{102}$$

From eqs. (100) and (101) we find

$$\tau_0 = t - \frac{r}{c} \tag{103}$$

where c is the speed of light.

Using eq. (103) in (94) we find

$$\Phi^a(x) = A_0^a(t, \mathbf{x}) = \frac{q^b(t - \frac{r}{c})}{r} \left[\frac{\exp[g \int dr \frac{Q(t-\frac{r}{c})}{r}] - 1}{g \int dr \frac{Q(t-\frac{r}{c})}{r}} \right]_{ab} \tag{104}$$

which reproduces eq. (12) where dr integration is an indefinite integration, $\vec{X}(\tau_0)$ is the position vector of the quark at the retarded time,

$$r = |\vec{x} - \vec{X}(\tau_0)|, \qquad Q_{ab}(\tau_0) = f^{abd} q^d(\tau_0) \tag{105}$$

and the repeated color indices b, d(=1,2,...8) are summed.

Hence we find from eq. (104) that, unlike Coulomb potential $A_0(\mathbf{x}) = \frac{e}{r}$ in Maxwell theory which is independent of time t, the color potential $A_0^a(t, \mathbf{x})$ produced by the color charges of the quark in Yang-Mills theory depends on time t even if $\vec{\beta} = 0$. This is a consequence of time dependent color charges of the quark in Yang-Mills theory. The color potential (Yang-Mills potential) $A_0^a(x)$ at time t depends on color charges of the quark at the retarded time $t - \frac{r}{c}$.

Note that when the color charge q^a is constant we find from eq. (104)

$$\Phi^a(x) = \frac{q^a}{r} \tag{106}$$

which reproduces the Coulomb-like potential, similar to Maxwell theory where the constant electric charge e produces Coulomb potential $\Phi(x) = \frac{e}{r}$.

The exact form of color potential produced by color charges of the quark may provide an insight to the question "why quarks are confined inside a (stable) proton". Hence in order to find the exact form of the color potential produced by the color charges of the quark we need to find the exact form of eight time dependent fundamental color charges $q^a(t)$ of the quark where $a = 1, 2, ...8$ are color indices. The derivation of general form of eight time dependent fundamental color charges $q^a(t)$ of a quark in Yang-Mills theory in SU(3) will be presented in a separate publication.

V. CONCLUSION

Constant electric charge e satisfies the continuity equation $\partial_\mu j^\mu(x) = 0$ where $j^\mu(x)$ is the current density of the electron. However, the Yang-Mills color current density $j^{\mu a}(x)$ of the quark satisfies the equation $D_\mu[A]j^{\mu a}(x) = 0$ which is not a continuity equation ($\partial_\mu j^{\mu a}(x) \neq 0$) which implies that a color charge $q^a(t)$ of the quark is not constant but it is time dependent where $a = 1, 2, ...8$ are color indices. In this paper we have derived general form of color potential produced by color charges of the quark. We have found that the general form of the color potential produced by the color charges of the quark at rest is given by $\Phi^a(x) = A_0^a(t, \mathbf{x}) = \frac{q^b(t-\frac{r}{c})}{r}\left[\frac{\exp[g\int dr \frac{Q(t-\frac{r}{c})}{r}]-1}{g\int dr \frac{Q(t-\frac{r}{c})}{r}}\right]_{ab}$ where dr integration is an indefinite integration, $Q_{ab}(\tau_0) = f^{abd}q^d(\tau_0)$, $r = |\vec{x} - \vec{X}(\tau_0)|$, $\tau_0 = t - \frac{r}{c}$ is the retarded time, c is the speed of light, $\vec{X}(\tau_0)$ is the position of the quark at the retarded time and the repeated color indices b, d(=1,2,...8) are summed. For constant color charge q^a we reproduce the Coulomb-like potential $\Phi^a(x) = \frac{q^a}{r}$ which is consistent with the Maxwell theory where constant electric charge e produces the Coulomb potential $\Phi(x) = \frac{e}{r}$.

Acknowledgments

I thank George Sterman for useful discussions, in particular, with respect to Lorentz transformation near speed of light. I also thank Robert Shrock for discussions on lattice

QCD.

Appendix A: Derivation of Abelian-Like Non-Abelian Pure Gauge Potential

The solution of the inhomogeneous wave equation

$$\partial_\nu \partial^\nu \mathcal{A}^a_\mu(x) = \mathcal{J}^a_\mu(x). \tag{A1}$$

is given by

$$\mathcal{A}^a_\mu(x) = \int d^4x' D_r(x-x') \mathcal{J}^a_\mu(x') \tag{A2}$$

where $D_r(x-x')$ is the retarded Greens function [9]. When $\mathcal{A}^a_\mu(x)$ satisfies the Lorentz gauge condition

$$\partial^\mu \mathcal{A}^a_\mu(x) = 0, \tag{A3}$$

then eq. (A1) reduces to the inhomogeneous Maxwell-like equation

$$\partial^\mu \mathcal{F}^a_{\mu\nu}(x) = \mathcal{J}^a_\nu(x) \tag{A4}$$

where

$$\mathcal{F}^a_{\mu\nu}(x) = \partial_\mu \mathcal{A}^a_\nu(x) - \partial_\nu \mathcal{A}^a_\mu(x). \tag{A5}$$

Note that $\mathcal{A}^a_\mu(x)$ in eq. (A1) is at any arbitrary four-velocity of the charge. Only when the four velocity of the charge is at speed of light we call $\mathcal{A}^{\mu a}(x)$ as the corresponding pure gauge potential.

Hence we find that if $\mathcal{A}^a_\mu(x)$ obtained from eq. (A2) satisfies Lorentz gauge condition eq. (A3) then it satisfies Maxwell-like eq. (A4).

1. Abelian-Like Potential Produced From Constant Color Charge

The color current density of the constant color charge q^a is given by [9]

$$\mathcal{J}^{\mu a}(x) = \int d\tau \; q^a \; u^\mu(\tau) \; \delta^{(4)}(x - X(\tau)) \tag{A6}$$

where $X^\mu(\tau)$ is the time-space position of the point charge q^a and

$$u^\mu(\tau) = \frac{dX^\mu(\tau)}{d\tau} \tag{A7}$$

is the four-velocity of the charge. Note that eq. (A6) satisfies the continuity equation

$$\partial^\mu \mathcal{J}^a_\mu(x) = 0 \tag{A8}$$

which confirms that the color charge q^a is constant.

Using eq. (A6) in (A2) we find that the abelian-like potential produced from constant color charge q^a is given by

$$\mathcal{A}^a_\mu(x) = q^a \frac{\beta_\mu(\tau_0)}{\beta(\tau_0)\cdot(x - X(\tau_0))} \tag{A9}$$

where τ_0 is determined from the solution of the retarded equation

$$x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)|. \tag{A10}$$

a. Abelian-Like Potential Satisfies Lorentz Gauge Condition

From eq. (A10) we find

$$\partial^\mu[u(\tau_0)\cdot(x - X(\tau_0))] = [\frac{[\dot{u}(\tau_0)\cdot(x - X(\tau_0)) - 1](x - X(\tau_0))^\mu}{u(\tau_0)\cdot(x - X(\tau_0))} + u^\mu(\tau_0)] \tag{A11}$$

where

$$\dot{u}^\mu(\tau_0) = \frac{du^\mu(\tau)}{d\tau}|_{\tau=\tau_0}. \tag{A12}$$

From eqs. (A9) and (A10) we find

$$\partial^\nu \mathcal{A}^a_\mu(x) = q^a \frac{(x - X(\tau_0))^\nu \dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x - X(\tau_0))]^2} - q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x - X(\tau_0))]^2}[\frac{[\dot{u}(\tau_0)\cdot(x - X(\tau_0)) - 1](x - X(\tau_0))^\nu}{u(\tau_0)\cdot(x - X(\tau_0))} + u^\nu(\tau_0)] \tag{A13}$$

which gives

$$\partial^\mu \mathcal{A}^a_\mu(x) = q^a \frac{(x - X(\tau_0))^\mu \dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x - X(\tau_0))]^2} - q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x - X(\tau_0))]^2}[\frac{[\dot{u}(\tau_0)\cdot(x - X(\tau_0)) - 1](x - X(\tau_0))^\mu}{u(\tau_0)\cdot(x - X(\tau_0))} + u^\mu(\tau_0)]. \tag{A14}$$

Hence we find from eq. (A14) that the abelian-like potential as given by eq. (A9) satisfies the Lorentz gauge condition

$$\partial^\mu \mathcal{A}^a_\mu(x) = 0. \tag{A15}$$

b. *Abelian-Like Potential Satisfies Maxwell-Like Equation*

From eq. (A13) and (A10) we find

$$\begin{aligned}
&\partial_\nu \partial^\nu \mathcal{A}^a_\mu(x) = q^a \frac{\partial_\nu[(x-X(\tau_0))^\nu \dot{u}_\mu(\tau_0)]}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} - 2q^a \frac{(x-X(\tau_0))^\nu \dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[u_\nu(\tau_0)] \\
&-q^a \frac{(x-X(\tau_0))_\nu \dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[u^\nu(\tau_0)] + 2q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[2\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1] \\
&-q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}[\frac{[[\dot{u}(\tau_0)\cdot\partial_\nu(x-X(\tau_0))]](x-X(\tau_0))^\nu}{u(\tau_0)\cdot(x-X(\tau_0))}] \\
&-3q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}[\frac{[\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1]}{u(\tau_0)\cdot(x-X(\tau_0))}] \\
&+q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1] - q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[\dot{u}(\tau_0)\cdot(x-X(\tau_0))]
\end{aligned} \tag{A16}$$

which gives

$$\begin{aligned}
&\partial_\nu \partial^\nu \mathcal{A}^a_\mu(x) = 3q^a \frac{\dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} - 2q^a \frac{\dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} - q^a \frac{\dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} \\
&+q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[\dot{u}(\tau_0)\cdot(x-X(\tau_0))+1] - q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[\dot{u}(\tau_0)\cdot(x-X(\tau_0))] \\
&+q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1] - q^a \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}[\dot{u}(\tau_0)\cdot(x-X(\tau_0))].
\end{aligned} \tag{A17}$$

Hence from eq. (A17) we find

$$\partial_\nu \partial^\nu \mathcal{A}^a_\mu(x) = 0. \tag{A18}$$

Eqs. (A18) and (A15) give

$$\partial^\mu \mathcal{F}^a_{\mu\nu}(x) = 0 \tag{A19}$$

where $\mathcal{F}^a_{\mu\nu}(x)$ is given by eq. (A5).

Since $x^\mu \neq X^\mu(\tau_0)$ is the observation point of the potential (or any gauge invariant obtained from the potential where $X^\mu(\tau_0)$ is the time-space position of the charge), we find from eqs. (A19) and (A6) that the abelian-like potential in eq. (A9) satisfies Maxwell-like eq. (A4).

c. *Abelian-Like Pure Gauge Potential*

Note that when the speed of the charge is arbitrarily close to the speed of light ($v \sim c$) we find from eq. (A9) that the abelian-like (approximate) pure gauge potential produced from the constant color charge q^a is given by

$$\mathcal{A}^a_\mu(x) = q^a \frac{\beta^\mu_{\sim c}}{\beta_{\sim c} \cdot (x - X(\tau_0))} \sim \partial_\mu \omega^a(x), \qquad \omega^a(x) = q^a \ \ln[\beta_{\sim}c \cdot (x - X(\tau_0))] \tag{A20}$$

which reproduces eq. (49) where $\beta^\mu_{\sim c}$ is given by eq. (40).

2. Abelian-Like Non-abelian Pure Gauge Potential Produced From Time Dependent Color Charge

Extending eq. (A6) to time dependent color charge we find that

$$\mathcal{J}^{\mu a}(x) = \int d\tau \ q^a(\tau) \ u^\mu(\tau) \ \delta^{(4)}(x - X(\tau)) \tag{A21}$$

where $u^\mu(\tau)$ is any arbitrary four velocity of the charge (not necessarily at speed of light). From eq. (A21) we find that

$$\partial^\mu \mathcal{J}^a_\mu(x) \ \neq \ 0 \tag{A22}$$

which is consistent with the fact that the color charge $q^a(\tau)$ is not constant.

Using eq. (A21) in (A2) we find that

$$\mathcal{A}^a_\mu(x) = q^a(\tau_0) \frac{u_\mu(\tau_0)}{u(\tau_0) \cdot (x - X(\tau_0))} \tag{A23}$$

where τ_0 is determined from the solution of the retarded equation

$$x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)|. \tag{A24}$$

From eqs. (A23) and (A24) we find

$$\partial^\nu \mathcal{A}^a_\mu(x) = q^a(\tau_0) \frac{(x - X(\tau_0))^\nu \dot{u}_\mu(\tau_0)}{[u(\tau_0) \cdot (x - X(\tau_0))]^2} - q^a(\tau_0) \frac{u_\mu(\tau_0)}{[u(\tau_0) \cdot (x - X(\tau_0))]^2}$$
$$[\frac{[\dot{u}(\tau_0) \cdot (x - X(\tau_0)) - 1](x - X(\tau_0))^\nu}{u(\tau_0) \cdot (x - X(\tau_0))} + u^\nu(\tau_0)] + [\partial^\nu q^a(\tau_0)] \frac{u_\mu(\tau_0)}{u(\tau_0) \cdot (x - X(\tau_0))}. \tag{A25}$$

From eq. (A25) we find

$$\begin{aligned}
&\partial_\nu \partial^\nu \mathcal{A}^a_\mu(x) = [\partial_\nu q^a(\tau_0)] \frac{(x-X(\tau_0))^\nu \dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} - [\partial_\nu q^a(\tau_0)] \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} \\
&[\frac{[\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1](x-X(\tau_0))^\nu}{u(\tau_0)\cdot(x-X(\tau_0))} + u^\nu(\tau_0)] + [\partial^\nu q^a(\tau_0)][\partial_\nu[\frac{u_\mu(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))}]] \\
&+[\partial_\nu\partial^\nu q^a(\tau_0)]\frac{u_\mu(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))}
\end{aligned} \tag{A26}$$

which gives

$$\begin{aligned}
&\partial_\nu \partial^\nu \mathcal{A}^a_\mu(x) = -\dot{q}^a(\tau_0) \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} - [\frac{(x-X(\tau_0))_\nu \dot{q}^a(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))}] \\
&[\frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}][\frac{[\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1](x-X(\tau_0))^\nu}{u(\tau_0)\cdot(x-X(\tau_0))} + u^\nu(\tau_0)] \\
&+[\frac{(x-X(\tau_0))^\nu (x-X(\tau_0))_\nu \ddot{q}^a(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}] \frac{u_\mu(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))} \\
&+[\frac{[\partial_\nu(x-X(\tau_0))^\nu]\dot{q}^a(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))}] \frac{u_\mu(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))} \\
&-[\frac{(x-X(\tau_0))^\nu \dot{q}^a(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}] \frac{u_\mu(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))} [\frac{[\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1](x-X(\tau_0))_\nu}{u(\tau_0)\cdot(x-X(\tau_0))} + u_\nu(\tau_0)]
\end{aligned} \tag{A27}$$

where

$$\dot{q}^a(\tau_0) = \frac{dq^a(\tau)}{d\tau}|_{\tau=\tau_0}. \tag{A28}$$

We find from eq. (A27)

$$\begin{aligned}
\partial_\nu \partial^\nu \mathcal{A}^a_\mu(x) = &-\dot{q}^a(\tau_0) \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} - \dot{q}^a(\tau_0) \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} \\
&+3\dot{q}^a(\tau_0) \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} - \dot{q}^a(\tau_0) \frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}
\end{aligned} \tag{A29}$$

which gives

$$\partial_\nu \partial^\nu \mathcal{A}^a_\mu(x) = 0. \tag{A30}$$

Since $x^\mu \neq X^\mu(\tau_0)$ is the observation point of the potential (or any gauge invariant obtained from the potential where $X^\mu(\tau_0)$ is the time-space position of the charge), we find that eq. (A30) is expected because eq. (A23) is obtained from eq. (A2) where $\mathcal{J}^a_\mu(x)$ is given by eq. (A21).

a. *Abelian-Like Non-Abelian Pure Gauge Potential Satisfies Lorentz Gauge Condition*

From eq. (A25) we find

$$\partial^\mu \mathcal{A}^a_\mu(x) = q^a(\tau_0)\frac{(x-X(\tau_0))^\mu \dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} - q^a(\tau_0)\frac{u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2} \\ [\frac{[\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1](x-X(\tau_0))^\mu}{u(\tau_0)\cdot(x-X(\tau_0))} + u^\mu(\tau_0)] + [\dot{q}^a(\tau_0)]\frac{(x-X(\tau_0))^\mu}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}u_\mu(\tau_0) \tag{A31}$$

which gives

$$\partial^\mu \mathcal{A}^a_\mu(x) = \frac{\dot{q}^a(\tau_0)}{u(\tau_0)\cdot(x-X(\tau_0))}, \qquad \text{where} \qquad \dot{q}^a(\tau_0) = \frac{dq^a(\tau)}{d\tau}|_{\tau=\tau_0}. \tag{A32}$$

For unform velocity we find from eq. (A32)

$$\partial^\mu \mathcal{A}^a_\mu(x) = \frac{\dot{q}^a(\tau_0)}{u\cdot(x-X(\tau_0))}. \tag{A33}$$

When the quark in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$, see eq. (40)) we find from eq. (A33) that the Lorentz gauge condition is (approximately) satisfied, *i.e.*,

$$\partial^\mu \mathcal{A}^a_\mu(x) \sim 0. \tag{A34}$$

b. *Abelian-Like Non-Abelian Pure Gauge Potential Satisfies Maxwell-Like Equation*

From eqs. (A23), (A34) and (A30) we find that the abelian-like non-abelian potential

$$\mathcal{A}^{\mu a}(x) = q^a(\tau_0)\frac{\beta^\mu_{\sim c}}{\beta_{\sim c}\cdot(x-X(\tau_0))} \tag{A35}$$

which is produced by the time dependent color charge of the quark in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) satisfies the equation

$$\partial^\nu \mathcal{F}^a_{\nu\mu}(x) \sim 0 \tag{A36}$$

where $\mathcal{F}^a_{\mu\nu}(x)$ is given by eq. (A5).

Since $x^\mu \neq X^\mu(\tau_0)$ is the observation point of the potential (or any gauge invariant obtained from the potential where $X^\mu(\tau_0)$ is the time-space position of the charge), we find that eq. (A36) satisfies Maxwell-like equation (A4) where the abelian-like non-abelian color current density $\mathcal{J}^{\mu a}(x)$ is given by eq. (A21) and $\mathcal{F}^a_{\mu\nu}(x)$ is given by eq. (A5).

c. Abelian-Like Non-Abelian Pure Gauge Potential Gives $\mathcal{F}^a_{\mu\nu}(x)=0$

From eqs. (A23) and (A24) we find

$$\partial_\mu \mathcal{A}^a_\nu(x) - \partial_\nu \mathcal{A}^a_\mu(x) = q^a(\tau_0)\frac{(x-X(\tau_0))_\mu \dot{u}_\nu(\tau_0) - (x-X(\tau_0))_\nu \dot{u}_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}$$
$$-q^a(\tau_0)\ [\frac{[\dot{u}(\tau_0)\cdot(x-X(\tau_0))-1][(x-X(\tau_0))_\mu u_\nu(\tau_0) - (x-X(\tau_0))_\nu u_\mu(\tau_0)]}{[u(\tau_0)\cdot(x-X(\tau_0))]^3}]$$
$$+[\dot{q}^a(\tau_0)]\frac{(x-X(\tau_0))_\mu u_\nu(\tau_0) - (x-X(\tau_0))_\nu u_\mu(\tau_0)}{[u(\tau_0)\cdot(x-X(\tau_0))]^2}. \tag{A37}$$

For uniform velocity we obtain from eq. (A37)

$$\partial_\mu \mathcal{A}^a_\nu(x) - \partial_\nu \mathcal{A}^a_\mu(x) = q^a(\tau_0)\ [\frac{(x-X(\tau_0)_\mu u_\nu - (x-X(\tau_0))_\nu u_\mu}{[u\cdot(x-X(\tau_0))]^3}]$$
$$+[\dot{q}^a(\tau_0)]\frac{(x-X(\tau_0))_\mu u_\nu - (x-X(\tau_0))_\nu u_\mu}{[u\cdot(x-X(\tau_0))]^2}. \tag{A38}$$

When the quark in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v\sim c$, see eq. (40), we find from eq. (A38)

$$\partial_\mu \mathcal{A}^a_\nu(x) - \partial_\nu \mathcal{A}^a_\mu(x) \sim 0. \tag{A39}$$

When the quark in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v\sim c$) we find from eqs. (A39) and (A35) that

$$\mathcal{F}^a_{\mu\nu}(x) \sim 0 \tag{A40}$$

where $\mathcal{F}^a_{\mu\nu}(x)$ is given by eq. (A5).

3. General Form of the Abelian-Like Non-Abelian Pure Gauge Potential Produced From Time Dependent Color Charge $q^a(\tau)$

From eqs. (A35), (A34), (A36) and (A40) we find that

$$\mathcal{A}^{\mu a}(x) = q^a(\tau_0)\frac{\beta^\mu_{\sim c}}{\beta_{\sim c}\cdot(x-X(\tau_0))}. \tag{A41}$$

satisfies

$$\partial^\mu \mathcal{A}^a_\mu(x) \sim 0, \tag{A42}$$

$$\partial^\mu \mathcal{F}^a_{\mu\nu}(x) \sim 0 \tag{A43}$$

and

$$\mathcal{F}^a_{\mu\nu}(x) \sim 0 \tag{A44}$$

where $\mathcal{F}^a_{\mu\nu}(x)$ is given by eq. (A5).

Hence we find that eq. (A41) is the general form of the abelian-like non-abelian (approximate) pure gauge potential produced from the time dependent color charge $q^a(t)$ of the quark in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$).

From eqs. (A41), (A44) and (A5) we find that eq. (A41) can be written as

$$\mathcal{A}^a_\mu(x) = q^a(\tau_0)\frac{\beta^\mu_{\sim c}}{\beta_{\sim c}\cdot(x-X(\tau_0))} \sim \partial_\mu \omega^a(x), \qquad \omega^a(x) = \int dl_c\, \frac{q^a(\tau_0)}{l_c}, \qquad l_c = \beta_{\sim c}\cdot(x-X(\tau_0)), \tag{A45}$$

which reproduce eq. (53).

Appendix B: Lorentz Transformation on Electromagnetic Field Produced by an Electron Moving Arbitrarily Close to the Speed of Light

The electromagnetic field produced by the electron of charge e moving with uniform velocity $\vec{v}$ is given by [9]

$$F^{\mu\nu}(x) = e\frac{(x-X(\tau_0))^\mu u^\nu - (x-X(\tau_0))^\nu u^\mu}{[u\cdot(x-X(\tau_0))]^3} \tag{B1}$$

where x^μ is the time-space position of the observation point of the electromagnetic field and $X^\mu(\tau)$ is the time-space position of the charge and

$$u^\mu = \frac{dX^\mu(\tau)}{d\tau} \tag{B2}$$

is the four-velocity of the charge. τ_0 is obtained from the solution of the retarded equation

$$x_0 - X_0(\tau_0) = |\vec{x} - \vec{X}(\tau_0)| = R. \tag{B3}$$

The Lorentz transformation of the electromagnetic field is given by [9]

$$F'^{\mu\nu} = \frac{\partial x'^\mu}{\partial x^\alpha}\frac{\partial x'^\nu}{\partial x^\beta}F^{\alpha\beta}(x), \qquad F'_{\mu\nu} = \frac{\partial x^\alpha}{\partial x'^\mu}\frac{\partial x^\beta}{\partial x'^\nu}F_{\alpha\beta}(x). \tag{B4}$$

For electron in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) we find from eqs. (B1), (41) and (40)

$$F^{\mu\nu}(x) = e\frac{(x - X(\tau_0))^\mu u^\nu_{\sim c} - (x - X(\tau_0))^\nu u^\mu_{\sim c}}{[u_{\sim c} \cdot (x - X(\tau_0))]^3}. \tag{B5}$$

From eqs. (B5), (41) and (40) we find

$$F^{\mu\nu}(x) \sim 0. \tag{B6}$$

Hence from eqs. (B5) and (B6) we find that when electron in uniform motion is at its highest speed (which is arbitrarily close to the speed of light $v \sim c$), it produces (approximate) pure gauge electromagnetic field. We call it (approximate) pure gauge electromagnetic field $F^{\mu\nu}(x)$ because electron has non-zero mass (even if very small) and hence it can not travel exactly at the speed of light $v = c$ to produce exact pure gauge electromagnetic field

$$F^{\mu\nu}(x) = 0. \tag{B7}$$

Hence for electron we find from eq. (B6)

$$F^{\mu\nu}(x) \neq 0, \qquad \text{when}, \qquad v \sim c, \qquad \text{even if} \qquad F^{\mu\nu}(x) \sim 0. \tag{B8}$$

Similarly from eqs. (42) and (B5) we find that

$$F^{\mu\nu}(x) \neq \infty, \qquad \text{when}, \qquad v \sim c. \tag{B9}$$

Since x'_0 and x'^i are not exactly infinity (see eq. (70)) and since $F^{\mu\nu}(x)$ in eq. (B6) (see eqs. (B5), (B8) and (B9)) is neither (exactly) infinity nor (exactly) zero we can perform Lorentz transformation on $F^{\mu\nu}(x)$ in eq. (B5).

From eq. (B4) we find

$$F'^{0i} = \frac{\partial x'_0}{\partial x^\alpha}\frac{\partial x'^i}{\partial x^\beta} F^{\alpha\beta}(x). \tag{B10}$$

From eqs. (B10) and (69) we find

$$F'^{0i} = u_{\alpha\sim c}[g_{i\delta} + \frac{\gamma_{\sim c} - 1}{\vec{\beta}^2_{\sim c}} \beta_{j\sim c} g_{j\delta} \beta^i_{\sim c} - \gamma_{\sim c} \beta^i_{\sim c} g_{0\delta}] F^{\alpha\delta}(x). \tag{B11}$$

Since $F^{\mu\nu}(x)$ in eq. (B5) is neither (exactly) infinity nor (exactly) zero (see eqs. (B9) and (B8)) we find by using eq. (B5) in (B11)

$$F'^{0i} = \frac{e}{[u_{\sim c}\cdot(x-X(\tau_0))]^3}\ \left[[g^i_\delta + \frac{\gamma_{\sim c}-1}{\vec{\beta}^2_{\sim c}}\beta_{j\sim c}g_{j\delta}\beta^i_{\sim c} - \gamma_{\sim c}\beta^i_{\sim c}g_{0\delta}]\right]$$
$$[u_{\sim c}\cdot(x-X(\tau_0))u^\delta_{\sim c} - (x-X(\tau_0))^\delta]. \tag{B12}$$

Since electron has non-zero mass we find from eq. (42) that $F'^{0i}\neq 0$ and $F'^{0i}\neq\infty$ in eq. (B12). Hence we find from eq. (B12)

$$F'^{0i} = \frac{e}{[u_{\sim c}\cdot(x-X(\tau_0))]^3}\ [[u_{\sim c}\cdot(x-X(\tau_0))u^i_{\sim c} - (x-X(\tau_0))^i] + [\frac{\gamma_{\sim c}-1}{\vec{\beta}^2_{\sim c}}\beta_{j\sim c}\beta^i_{\sim c}]$$
$$[u_{\sim c}\cdot(x-X(\tau_0))u_{j\sim c} - (x-X(\tau_0))_j] - \gamma_{\sim c}\beta^i_{\sim c}[u_{\sim c}\cdot(x-X(\tau_0))u_{0\sim c} - (x-X(\tau_0))_0]]. \tag{B13}$$

Simplifying eq. (B13) we find

$$F'^{0i} = \frac{e}{[u_{\sim c}\cdot(x-X(\tau_0))]^3}\ [[u_{\sim c}\cdot(x-X(\tau_0))u^i_{\sim c} - (x-X(\tau_0))^i]$$
$$+(\gamma_{\sim c}-1)\beta^i_{\sim c}[u_{\sim c}\cdot(x-X(\tau_0))\gamma_{\sim c} - \frac{\vec{\beta}_{\sim c}\cdot(\vec{x}-\vec{X}(\tau_0))}{\vec{\beta}^2_{\sim c}}]$$
$$-\gamma_{\sim c}\beta^i_{\sim c}[u_{\sim c}\cdot(x-X(\tau_0))\gamma_{\sim c} - (x_0-X_0(\tau_0))]]. \tag{B14}$$

Using eqs. (69) and (73) in (B14) we find

$$F'^{0i} = \frac{e}{(x'_0-X'_0(\tau'_0))^3}\ [[(x'_0-X'_0(\tau'_0)u^i_{\sim c} - (x-X(\tau_0))^i]$$
$$+(\gamma_{\sim c}-1)\beta^i_{\sim c}[(x'_0-X'_0(\tau'_0))\gamma_{\sim c} - \frac{\vec{\beta}_{\sim c}\cdot(\vec{x}-\vec{X}(\tau_0))}{\vec{\beta}^2_{\sim c}}] - \gamma_{\sim c}\beta^i_{\sim c}[(x'_0-X'_0(\tau_0))\gamma_{\sim c} - (x_0-X_0(\tau_0))]]. \tag{B15}$$

The retarded condition is given by

$$x'_0 - X'_0(\tau_0) = |\vec{x}' - \vec{X}'(\tau'_0)| = R' \tag{B16}$$

By using eq. (B16) in (B15) we find

$$F'^{0i} = \frac{e}{R'^3}\ \left[R'\gamma_{\sim c}\beta^i_{\sim c} - Rn^i + (\gamma_{\sim c}-1)\beta^i_{\sim c}[R'\gamma_{\sim c} - R\frac{\vec{\beta}_{\sim c}\cdot\vec{n}}{\vec{\beta}^2_{\sim c}}] - \gamma_{\sim c}\beta^i_{\sim c}[R'\gamma_{\sim c} - R]\right] \tag{B17}$$

where we have introduced the notation

$$\vec{R} = \vec{n}R = \vec{x} - \vec{X}(\tau_0). \tag{B18}$$

From eq. (B17) we find

$$F'^{0i} = \frac{eR}{R'^3}\left[-n^i + \frac{\vec{\beta}_{\sim c}\cdot\vec{n}}{\vec{\beta}^2_{\sim c}}\beta^i_{\sim c} + \gamma_{\sim c}\beta^i_{\sim c}[1 - \frac{\vec{\beta}_{\sim c}\cdot\vec{n}}{\vec{\beta}^2_{\sim c}}]\right] \tag{B19}$$

which gives

$$F'^{0i} = -\frac{e}{R'^3}\left[R^i - \frac{\vec{\beta}_{\sim c}\cdot\vec{R}}{\vec{\beta}^2_{\sim c}}\beta^i_{\sim c} - \gamma_{\sim c}\beta^i_{\sim c}[R - \frac{\vec{\beta}_{\sim c}\cdot\vec{R}}{\vec{\beta}^2_{\sim c}}]\right]. \tag{B20}$$

Eq. (B20) can be simplified to find

$$F'^{0i} = -\frac{e}{R'^3}\left[R^i + (\gamma_{\sim c} - 1)\frac{\vec{\beta}_{\sim c}\cdot\vec{R}}{\vec{\beta}^2_{\sim c}}\beta^i_{\sim c} - \gamma_{\sim c}\beta^i_{\sim c}(x_0 - X_0(\tau_0))\right]. \tag{B21}$$

Hence we find from eqs. (B16) and (B21)

$$F'^{0i} = -\frac{e}{|\vec{x}' - \vec{X}'(\tau'_0)|^3}\left[(x^i - X^i(\tau_0)) + (\gamma_{\sim c} - 1)\frac{\vec{\beta}_{\sim c}\cdot(\vec{x} - \vec{X}(\tau_0))}{\vec{\beta}^2_{\sim c}}\beta^i_{\sim c} - \gamma_{\sim c}\beta^i_{\sim c}(x_0 - X_0(\tau_0))\right]. \tag{B22}$$

By using eqs. (69) and (73) in eq. (B22) we find

$$F'^{0i} = -e\frac{x'^i - X'^i(\tau'_0)}{|\vec{x}' - \vec{X}'(\tau'_0)|^3}. \tag{B23}$$

Hence from eqs. (73) and (B23) we find

$$F^{0i} = -e\frac{x^i - X^i(\tau_0)}{|\vec{x} - \vec{X}(\tau_0)|^3} \tag{B24}$$

which is the exact form of the Coulomb electric field produced by the electron at rest.

Hence we find that the exact form of the Coulomb electric field in eq. (B24) produced by the electron at rest can be obtained from the form of (approximate) pure gauge electromagnetic field $F^{\mu\nu}(x)$ in eq. (B5) [see eqs. (B6), (B8) and (B9)] produced by the electron in uniform motion at its highest speed (which is arbitrarily close to the speed of light $v \sim c$) by using Lorentz transformations.

Hence one finds that if one knows the form of the (approximate) pure gauge potential produced by a point charge of non-zero mass in uniform motion at its highest speed (which is arbitrarily close to the speed of the light $v \sim c$), then one can obtain the exact form of the potential produced by the same charge at rest by using Lorentz transformations. This is consistent with the replacement given by eq. (62) which can be used to obtain eq. (58) from (60) and eq. (B24) from (B5).

Appendix C: Yang-Mills Color Current Density

Note that the Lienard-Wiechert potential in eq. (35) can be obtained from the current density

$$j_\mu(x) = e \int d\tau \ u_\mu(\tau) \ \delta^{(4)}(x - X(\tau)) \tag{C1}$$

of the point charge e in Maxwell theory. However, the Yang-Mills potential in eq. (96) gives the Yang-Mills color current density

$$j^{\nu a}(x) = D_\mu[A] F^{\mu\nu a}(x) \tag{C2}$$

which is non-linear function of color charges $q^a(\tau)$. This implies that Yang-Mills color density

$$j^a_\mu(x) \neq \int d\tau \ u_\mu(\tau) \ q^a(\tau) \ \delta^{(4)}(x - X(\tau)). \tag{C3}$$

Hence one finds that not only the Yang-Mills potential $A^a_\mu(x)$ in eq. (96) is non-linear function of time dependent color charges $q^a(\tau)$ but the Yang-Mills color current density $j^a_\mu(x)$ in eq. (C2) is also non-linear function of color charges $q^a(\tau)$.

This can also be seen as follows. From eq. (4) one finds in Maxwell theory [10]

$$\int d^3x j^0(x) = e \int d^3x \psi^\dagger(x)\psi(x) = e \tag{C4}$$

and

$$\int d^3x \vec{j}(x) = e \int d^3x \bar{\psi}(x)\vec{\gamma}\psi(x) = e < \frac{\vec{p}}{m} > . \tag{C5}$$

Eq. (C1) is consistent with eqs. (C4) and (C5).

On the other hand from eqs. (11), (10), (9), (C2) and (96) one finds

$$\int d^3x j^a_0(x) = g \int d^3x \psi^\dagger_i(x) T^a_{ij} \psi_j(x) \neq q^a(t) \tag{C6}$$

and

$$\int d^3x \vec{j}^a(x) = g \int d^3x \bar{\psi}_i(x) \vec{\gamma} T^a_{ij} \psi_j(x) \neq < \frac{\vec{p}}{m} > q^a(t). \tag{C7}$$

Eq. (C3) is consistent with eqs. (C6) and (C7).

---

[1] D. J. Gross and F. Wilczek, Phys.Rev.Lett.30:1343-1346,1973; Phys.Rev.D8:3633-3652,1973; D. Politzer, Phys.Rept.14:129-180,1974.

[2] K. G. Wilson, Phys. Rev. D10 (1974) 2445.

[3] C. N. Yang and R. L. Mills, Phys. Rev. 96 (1954) 191.

[4] see for example, T. Muta, *Foundations of Quantum Chromodynamics* , World Scientific Lecture Notes in Physics - Vol. 5.

[5] T. T. Wu and C. N. Yang, Phys. Rev. D12, 3843 (1975); *ibid* Phys. Rev. D13, 3233 (1976).

[6] R. Roskies, Phys. Rev. D15, 1722, 1731 (1977); J. Anandan and R. Roskies, Phys. Rev. D18, 1152 (1978); J. Math. Phys. 19, 2614 (1978).

[7] J. E. Mandula, Phys. Rev. D14, 3496 (1976); L. L. Wang and C. N. Yang, Phys. Rev. D17, 2687 (1978); L. S. Brown and D. B. Creamer, Phys. Rev. D18, 3695 (1978); M. Carmeli and M. Fischler, Phys. Rev. D19, 3653 (1979); P. Sikivie and N. Weiss, Phys. Rev. D18, 3809 (1978); *ibid* D20, 487 (1979); R. Jackiw, L. Jacobs and C. Rebbi, Phys. Rev. D20, 474 (1979); A. A. Actor, Rev. Mod. Phys. 51, 461 (1979); G. C. Nayak and R. Shrock, Phys. Rev. D77:045008, 2008; G. C. Nayak and P. Nieuwenhuizen, Phys.Rev.D71:125001,2005; Q. Wang *et al.*, Int.J.Mod.Phys.E10:483-500,2001; D. Dietrich, G. C. Nayak, W. Greiner, Phys.Rev.D64:074006,2001; D. Dietrich, G. C. Nayak, W. Greiner, J.Phys.G28:2001-2004,2002; D. Dietrich, G. C. Nayak, W. Greiner, hep-ph/0009178; G. C. Nayak, W. Greiner, hep-th/0001009; G. C. Nayak, D. Dietrich, W. Greiner, hep-ph/0104030; G. C. Nayak, Phys.Rev.D72:125010,2005; G. C. Nayak, Int.J.Mod.Phys.A25:3885-3898,2010.

[8] S. L. Adler, Phys.Rev.D17:3212,1978; P. Pirila, P. Presnajder, Nucl.Phys.B142:229,1978; R. A. Brandt, F. Neri, Nucl.Phys.B161:253-282,1979; A. Yildiz, P. H. Cox, Phys.Rev.D21:1095,1980; R. Jackiw, P. Rossi, Phys.Rev.D21:426,1980; R. A. Freedman, L. Wilets, S. D. Ellis, E. M. Henley, Phys.Rev.D22:3128,1980; C.H. Oh, R. Teh, W.K. Koo, Phys.Rev.D24:2305,1981; C.H. Oh, Phys.Rev.D25:2194,1982; G. C. Nayak, Eur.Phys.J.C64:73-79,2009; G. C. Nayak,

Int.J.Mod.Phys.A25:1155-1163,2010; G. C. Nayak, Eur.Phys.J.C59:715-722,2009; G. C. Nayak, Annals Phys.325:514-518,2010; G. C. Nayak, Electron.J.Theor.Phys.8:279-286,2011; G. C. Nayak, JHEP 0903:051,2009; J. S. Ball, Nucl.Phys.B201:352,1982; D. Horvat, Phys.Rev.D34:1197-1199,1986; I. R. Klebanov, J. M. Maldacena, JHEP 0604:024,2006.

[9] J. D. Jackson, *Classical Electrodynamics*, Second Edition, John Wiley and Sons.

[10] C. Itzykson and J-B. Zuber, *Quantum Field Theory*, Dover Publications Inc., Mineola, New York.